

JUNE 2017 CSEC MATHEMATICS PAPER 2

SECTION 1

1. (a) Using a calculator, or otherwise, calculate the EXACT value of

(i) $\left(4\frac{1}{3} - 1\frac{2}{5}\right) \div \frac{4}{15}$

SOLUTION:

Required to calculate: $\left(4\frac{1}{3} - 1\frac{2}{5}\right) \div \frac{4}{15}$ in exact form

Calculation:

$$\begin{aligned} 4\frac{1}{3} - 1\frac{2}{5} &= \frac{13}{3} - \frac{7}{5} \\ &= \frac{5(13) - 3(7)}{15} \\ &= \frac{65 - 21}{15} \\ &= \frac{44}{15} \end{aligned}$$

So, now we have

$$\begin{aligned} \left(4\frac{1}{3} - 1\frac{2}{5}\right) \div \frac{4}{15} &= \frac{44}{15} \div \frac{4}{15} \\ &= \frac{\cancel{44}^{11}}{\cancel{15}} \times \frac{\cancel{15}}{\cancel{4}_1} \\ &= 11 \text{ (in exact form)} \end{aligned}$$

(ii) $\frac{(3.1 - 1.15)^2}{0.005}$

SOLUTION:

Required to calculate: $\frac{(3.1 - 1.15)^2}{0.005}$ in exact form

Calculation:

$$\begin{aligned} \frac{(3.1 - 1.15)^2}{0.005} &= \frac{(1.95)^2}{0.005} \quad \text{(By calculator)} \\ &= \frac{3.8025}{0.005} \\ &= 760.5 \text{ (in exact form)} \end{aligned}$$

- (b) A store is promoting a new mobile phone under two plans. Plan A and Plan B. The plans are advertised as shown in the table below.

	Plan A	Plan B
Deposit	\$400	\$600
Monthly installment	\$65	\$80
Number of months to repay	12	6
Tax on ALL payments	0%	5%

- (i) Calculate the TOTAL cost of a phone under Plan A.

SOLUTION:

Data: Table showing two mobile phone plans as advertised by a store.

Required to calculate: The total cost of a phone under Plan A

Calculation:

Total cost under Plan A

$$\begin{aligned}
 &= \text{The deposit} + (\text{Monthly installments} \times \text{No. of months to repay}) + \text{Tax} \\
 &= \$400 + (\$65 \times 12) + \$0 \\
 &= \$(400 + 780) \\
 &= \$1180
 \end{aligned}$$

- (ii) Determine which of the two plans, A or B, is the better deal. Justify your answer.

SOLUTION:

Required to find: The better deal of the two plans

Solution:

Total cost under Plan B

$$\begin{aligned}
 &= \text{The deposit} + (\text{Monthly installments} \times \text{No. of months to repay}) + \text{Tax} \\
 &= \$600 + (\$80 \times 6) + \frac{5}{100}(\$600 + (\$80 \times 6)) \\
 &= \$(600 + 480) + \frac{5}{100}(\$600 + \$480) \\
 &= 1.05 \times \$(600 + 480) \\
 &= \$1134
 \end{aligned}$$

If the better deal is supposed to mean the plan that has a lesser cost, then Plan B is the better deal as there is a savings of $\$1180 - \$1134 = \$46$.

- (c) John's monthly electricity bill is based on the number of kWh of electricity that he consumes for that month. He is charged \$5.10 per kWh of electricity consumed. For the month of March 2016, two meter readings are displayed in the table below.

	Meter Readings (kWh)
Beginning 01 March	0 3 0 1 1
Ending 31 March	0 3 3 0 7

- (i) Calculate the TOTAL amount that John pays for electricity consumption for the month of March 2016.

SOLUTION:

Data: Table showing John's electricity meter readings, in kWh, at the beginning and end of March 2016. John is charged \$5.10 per kWh for electricity used.

Required to calculate: The total amount John pays for electricity for March 2016

Calculation:

Number of kWh used

= End reading on 31 March – Beginning reading on 01 March

$$\begin{array}{r}
 = 0\ 3\ 3\ 0\ 7 - \\
 \underline{0\ 3\ 0\ 1\ 1} \\
 2\ 9\ 6
 \end{array}$$

The cost at \$5.10 per kWh = $296 \times \$5.10 = \1509.60

∴ If John pays the full amount that is required, then he would pay \$1 509.60.

- (ii) For the next month, April 2016, John pays \$2 351.10 for electricity consumption. Determine his meter reading at the end of April 2016.

SOLUTION:

Data: John pays \$2 351.10 for electricity in April 2016.

Required to calculate: John's meter reading at the end of April 2016

Calculation:

For the month of April John pays \$2 351.10.

$$\begin{aligned}
 \text{At } \$5.10 \text{ per kWh, the number of kWh used in April} &= \frac{\$2351.10}{\$5.10} \\
 &= 461 \text{ kWh}
 \end{aligned}$$

Hence, the meter reading at the end of April should be the reading at the end of March + 461

$$\begin{array}{r}
 = 0\ 3\ 3\ 0\ 7\ + \\
 \quad \quad \quad 4\ 6\ 1 \\
 \hline
 0\ 3\ 7\ 6\ 8
 \end{array}$$

2. (a) Factorise the following expressions completely.

(i) $6y^2 - 18xy$

SOLUTION:

Required to factorise: $6y^2 - 18xy$

Solution:

$$\begin{aligned}
 6y^2 - 18xy &= 6y \times y - 6y \times 3x \\
 &= 6y(y - 3x)
 \end{aligned}$$

(ii) $4m^2 - 1$

SOLUTION:

Required to factorise: $4m^2 - 1$

Solution:

$$4m^2 - 1 = (2m)^2 - (1)^2$$

This is now in the form of the difference of two squares.

$$\text{So, } 4m^2 - 1 = (2m - 1)(2m + 1)$$

(iii) $2t^2 - 3t - 2$

SOLUTION:

Required to factorise: $2t^2 - 3t - 2$

Solution:

$$2t^2 - 3t - 2 = (2t + 1)(t - 2)$$

$$\left[\begin{array}{l} 2t^2 - 4t + t - 2 \\ 2t^2 - 3t - 2 \end{array} \right] \text{We may even expand to confirm our result}$$

(b) Write as a single fraction and simplify

$$\frac{5p+2}{3} - \frac{3p-1}{4}$$

SOLUTION:

Required to write: $\frac{5p+2}{3} - \frac{3p-1}{4}$ as a simplified single fraction.

Solution:

$$\begin{aligned} & \frac{5p+2}{3} - \frac{3p-1}{4} \\ & \frac{4(5p+2) - 3(3p-1)}{12} = \frac{20p+8-9p+3}{12} \\ & = \frac{20p-9p+8+3}{12} \\ & = \frac{11p+11}{12} \\ & = \frac{11(p+1)}{12} \end{aligned}$$

(c) A formula is given as $d = \sqrt{\frac{4h}{5}}$.

(i) Determine the value of d when $h = 29$. Give your answer correct to 3 significant figures.

SOLUTION:

Data: $d = \sqrt{\frac{4h}{5}}$ and $h = 29$

Required to calculate: d

Calculation:

When $h = 29$

$$d = \sqrt{\frac{4(29)}{5}}$$

$$= \sqrt{\frac{116}{5}}$$

$$= \sqrt{23.2}$$

$$= 4.816$$

$$= 4.82 \text{ (correct to 3 significant figures)}$$

(using the calculator)

(ii) Make h the subject of the formula.

SOLUTION:

Required to make: h the subject of the formula

Solution:

$$d = \sqrt{\frac{4h}{5}}$$

Squaring both sides to remove the hindrance of the root sign

$$d^2 = \frac{4h}{5}$$

$$\therefore 5 \times d^2 = 4h$$

$$4h = 5d^2$$

$$h = \frac{5d^2}{4}$$

3. (a) The Universal set, U , is defined as follows:

$$U = \{x : x \in N, 2 < x < 12\}$$

The sets M and R are subsets of U such that

$$M = \{\text{odd numbers}\}$$

$$R = \{\text{square numbers}\}$$

- (i) List the members of the subset M .

SOLUTION:

Data: $U = \{x : x \in N, 2 < x < 12\}$ and M and R are subsets of U where

$M = \{\text{odd numbers}\}$ and $R = \{\text{square numbers}\}$.

Required to list: The members of the subset M .

Solution:

$$U = \{3, 4, 5, 6, 7, 8, 9, 10, 11\}$$

$$\therefore M = \{3, 5, 7, 9, 11\}$$

- (ii) List the members of the subset R .

SOLUTION:

Required to list: The members of the subset R

Solution:


$$R = \{4, 9\}$$

- (iii) Draw a Venn diagram that represents the relationship among the defined subsets of U .

SOLUTION:

Required to draw: A Venn diagram to illustrate the information given about the sets U , M and R

Solution:




(b) Using a ruler, a pencil and a pair of compasses,



(i) construct accurately, the square $ABCD$, with sides 6 cm.


SOLUTION:

Required to construct: A square $ABCD$ with sides 6 cm.

Solution:








- (ii) construct, as an extension of your diagram in (b) (i), the trapezium $DABQ$ so that $\angle ABQ = 120^\circ$.

[Note: Credit will be given for clearly drawn construction lines.]

SOLUTION:

Required To Construct: An extension of the diagram previously down trapezium $DABQ$ with the $\angle ABQ = 120^\circ$.

Solution:



- (iii) Hence, measure and state the length of BQ .

SOLUTION:

Required to state: The length of BQ , by measurement

Solution:

$BQ = 6.9$ cm (by measurement with a ruler)

4. (a) The function f is defined as $f(x) = \frac{1}{3}x - 2$.

- (i) Find the value of $f(3) + f(-3)$.

SOLUTION:

Data: $f(x) = \frac{1}{3}x - 2$

Required to find: $f(3) + f(-3)$

Solution:

$$\begin{aligned} f(3) &= \frac{1}{3}(3) - 2 & f(-3) &= \frac{1}{3}(-3) - 2 \\ &= 1 - 2 & &= -1 - 2 \\ &= -1 & &= -3 \end{aligned}$$

So,

$$\begin{aligned} f(3) + f(-3) &= -1 + (-3) \\ &= -1 - 3 \\ &= -4 \end{aligned}$$

- (ii) Calculate the value of x for which $f(x) = 5$.

SOLUTION:

Required to calculate: x when $f(x) = 5$

Calculation:

$$\begin{aligned} f(x) &= 5 \\ \therefore \frac{1}{3}x - 2 &= 5 \\ \frac{1}{3}x &= 5 + 2 = 7 \\ x &= 7 \times 3 \quad (\times 3) \\ x &= 21 \end{aligned}$$

- (iii) Determine the inverse function, $f^{-1}(x)$

SOLUTION:

Required to find: $f^{-1}(x)$

Solution:

$$f(x) = \frac{1}{3}x - 2$$

$$\text{Let } y = \frac{1}{3}x - 2$$

Make x the subject of the formula:

$$y + 2 = \frac{1}{3}x$$


$$3(y + 2) = x$$

$$x = 3(y + 2)$$

Replace y by x to obtain:

$$f^{-1}(x) = 3(x + 2)$$

- (b) The graph below shows two straight lines, ℓ_1 and ℓ_2 . Line ℓ_1 intercepts the y -axis at $(0, 1)$. Line ℓ_2 intercepts the x and y axes at $(12, 0)$ and $(0, 6)$ respectively.



- (i) Calculate the gradient of the lines ℓ_1 and ℓ_2 .

SOLUTION:

Data: Diagram showing two lines ℓ_1 and ℓ_2 . The y – intercept of ℓ_1 is $(0, 1)$. The y – intercept of ℓ_2 is $(0, 6)$ and the x – intercept is $(12, 0)$.

Required to calculate: The gradient of ℓ_1 and ℓ_2

Calculation:

Two points on ℓ_1 are $(0, 1)$ and $(2, 5)$ which is the point of intersection of ℓ_1 and ℓ_2 . Each was obtained by a read-off.

$$\begin{aligned}\therefore \text{Gradient of } \ell_1 &= \frac{5 - 1}{2 - 0} \\ &= \frac{4}{2} \\ &= 2\end{aligned}$$

Two points on ℓ_2 are $(12, 0)$ and $(0, 6)$.

$$\begin{aligned}\therefore \text{Gradient of } \ell_2 &= \frac{6 - 0}{0 - 12} \\ &= -\frac{1}{2}\end{aligned}$$

- (ii) Determine the equation of the line ℓ_1

SOLUTION:

Required to find: The equation of the line ℓ_1

Solution:

The general equation of a straight line is of the form $y = mx + c$, where m is the gradient and c is the intercept on the y – axis. In this case, we have already found $m = 2$ and noted that $c = 1$.

\therefore The equation of the line ℓ_1 is $y = 2x + 1$.

- (iii) What is the relationship between ℓ_1 and ℓ_2 ? Give a reason for your answer.

SOLUTION:

Required to find: The relationship between ℓ_1 and ℓ_2 .

Solution:


$$\text{Gradient of } \ell_1 = 2$$

$$\text{Gradient of } \ell_2 = -\frac{1}{2}$$

$$\text{Gradient of } \ell_1 \times \text{Gradient of } \ell_2 = 2 \times -\frac{1}{2} = -1$$

Hence, ℓ_1 is perpendicular to ℓ_2 since the product of the gradients of perpendicular lines is -1 .

5. (a) $PTRS$, not drawn to scale, is a quadrilateral. Q is a point on PT such that $QT = QR = QP$. Angle $QRT = 76^\circ$.



Determine, given a reason for each step of your answer, the measure of

- (i) angle RQT


SOLUTION:

Data: Diagram showing a quadrilateral $PTRS$ such that $QT = QR = QP$ and angle $QRT = 76^\circ$.

Required to find:

\hat{RQT}

Solution:



Angle $RTQ = 76^\circ$

(The base angles of the isosceles triangle RQT are equal)

$$\therefore \text{Angle } RQT = 180^\circ - (76^\circ + 76^\circ)$$


$$= 28^\circ \quad (\text{The sum of interior angles in a triangle} = 180^\circ)$$

- (ii) angle PRT

SOLUTION:

Required to find: angle PRT

Solution:



Angle $QPR = \text{Angle } QRP$
(The base angles of the isosceles triangle RQP are equal)

Angle $QPR + \text{Angle } QRP = 28^\circ$
(The exterior angle of a triangle is equal to the sum of the interior opposite angles)

$$\therefore \text{Angle } QRP = \frac{28^\circ}{2} = 14^\circ$$

$$\text{Hence, angle } PRT = 76^\circ + 14^\circ = 90^\circ$$


(iii) angle SPT , given that angle $SRT = 145^\circ$ and angle $PSR = 100^\circ$.

SOLUTION:

Data: Angle $SRT = 145^\circ$ and angle $PSR = 100^\circ$

Required to find: angle SPT

Solution:




$$\begin{aligned} \text{Angle } SRP &= 145^\circ - 90^\circ \\ &= 55^\circ \end{aligned}$$

$$\begin{aligned} \text{Angle } SPR &= 180^\circ - (100^\circ + 55^\circ) \\ &= 25^\circ \quad (\text{Sum of the interior angles in a triangle} = 180^\circ) \end{aligned}$$

$$\begin{aligned} \text{Hence, angle } SPT &= 25^\circ + 14^\circ \\ &= 39^\circ \end{aligned}$$

- (b) The diagram below shows a triangle ABC and its image, $A'B'C'$, under a single transformation.




- (i) Describe completely the transformation that maps $\triangle ABC$ to $\triangle A'B'C'$.

SOLUTION:

Data: Diagram showing triangle ABC and its image $A'B'C'$ after a single transformation.

Required To Describe: The transformation that maps $\triangle ABC$ unto $\triangle A'B'C'$.

Solution:



$A'B'C'$ is congruent to ABC and is re-oriented. Hence, the transformation is a rotation. The perpendicular bisectors of BB' and AA' meet at O . Hence, O is the center of rotation. $\widehat{BOB'} = 90^\circ$ and the 'sweep' from OB to OB' is clockwise. The transformation is therefore a 90° clockwise rotation about O .

- (ii) The translation vector $T = \begin{pmatrix} 4 \\ -5 \end{pmatrix}$ maps $\Delta A'B'C'$ to $\Delta A''B''C''$. On the diagram above, draw the $\Delta A''B''C''$.


SOLUTION:

Data: $\Delta A'B'C'$ is mapped to $\Delta A''B''C''$ by $T = \begin{pmatrix} 4 \\ -5 \end{pmatrix}$.

Required To Draw: $\Delta A''B''C''$

Solution:

See graph, $A' = (1, 4)$, $B' = (1, 1)$ and $C' = (3, 1)$



$$\Delta A'B'C' \xrightarrow{T = \begin{pmatrix} 4 \\ -5 \end{pmatrix}} \Delta A''B''C''$$


$$A' (1, 4) \xrightarrow{\begin{pmatrix} 4 \\ -5 \end{pmatrix}} A'' (5, -1)$$

$$B' (1, 1) \xrightarrow{\begin{pmatrix} 4 \\ -5 \end{pmatrix}} B'' (5, -4)$$

$$C' (3, 1) \xrightarrow{\begin{pmatrix} 4 \\ -5 \end{pmatrix}} C'' (7, -4)$$

6. (a) In this problem take π to be $\frac{22}{7}$.

The diagram below, **not drawn to scale**, shows a field in the shape of a sector of a circle with center O and diameter 28 m. Angle POZ is 90° .



Calculate:

- (i) the area of the field

SOLUTION:

Data: Diagram showing a field POZ in the shape of a sector of a circle with center O and diameter 28 m. Angle POZ is 90° .

Required to calculate: The area of the field

Calculation:

$$\text{Radius} = \frac{\text{Diameter}}{2} = \frac{28 \text{ m}}{2} = 14 \text{ m}$$

$$\begin{aligned} \text{Area of the field} &= \frac{90^\circ}{360^\circ} \times \pi (14)^2 \\ &= \frac{1}{4} \times \frac{22}{7} \times 14 \times 14 \\ &= 154 \text{ m}^2 \end{aligned}$$

- (ii) the perimeter of the field

SOLUTION:

Required to calculate: The perimeter of the field

Calculation:

The perimeter of the field


= Length of radius OP + Length of arc PZ + Length of radius ZO

$$= 14 + \frac{90^\circ}{360^\circ} (2 \times \pi \times 14) + 14$$

$$= 14 + \left(\frac{1}{4} \times 2 \times \frac{22}{7} \times 14 \right) + 14$$

$$= 50 \text{ m}$$

- (b) The diagram below, **not drawn to scale**, shows a triangular prism $ABCDEF$. The cross-section is the right-angled triangle, ABC , where $AB = 6$ cm and $BC = 10$ cm.



Calculate


- (i) the area of the triangle ABC

SOLUTION:

Data: Diagram showing the triangular prism $ABCDEF$ which cross-section is a right-angled triangle. $AB = 6$ cm and $BC = 10$ cm.

Required to calculate: The area of triangle ABC

Calculation:



$$BC^2 = AC^2 + AB^2 \quad (\text{Pythagoras' Theorem})$$

$$(10)^2 = AC^2 + (6)^2$$

$$AC^2 = 100 - 36$$

$$AC^2 = 64$$

$$AC = \sqrt{64}$$

$$= 8 \text{ cm (taking the positive value of the root)}$$

$$\begin{aligned} \text{Area of } \triangle ABC &= \frac{6 \times 8}{2} \text{ cm}^2 \\ &= \frac{48}{2} \text{ cm}^2 \\ &= 24 \text{ cm}^2 \end{aligned}$$

- (ii) the length of the prism, if the volume is 540 cm^3

SOLUTION:

Data: The volume of the prism is 540 cm^3 .

Required to calculate: The length of the prism.

Calculation:

Volume of prism = Area of cross-section \times Length of prism

$$\therefore 540 = 24 \times \text{Length of prism}$$

$$\begin{aligned} \text{Length of prism} &= \frac{540}{24} \\ &= 22.5 \text{ cm} \end{aligned}$$

- (iii) the surface area of the prism.

SOLUTION:

Required to calculate: The surface area of the prism.

Calculation:

Surface area of the prism = Sum of the area of all five sides

$$\text{Area of } \triangle ABC = \frac{6 \times 8}{2} = 24 \text{ cm}^2$$

$$\text{Area of } \triangle DEF = \frac{6 \times 8}{2} = 24 \text{ cm}^2$$

$$\text{Area of rectangle } ABED = 6 \times 22.5 = 135 \text{ cm}^2$$

$$\text{Area of rectangle } ADFE = 8 \times 22.5 = 180 \text{ cm}^2$$

$$\text{Area of rectangle } BEFC = 10 \times 22.5 = 225 \text{ cm}^2$$

$$\begin{aligned} \text{Hence, the surface area of the prism} &= 24 + 24 + 135 + 180 + 225 \\ &= 588 \text{ cm}^2 \end{aligned}$$

7. The table below shows the speeds, to the nearest kmh^{-1} , of 90 vehicles that pass a checkpoint.

Speed (in kmh^{-1})	Frequency	Cumulative Frequency
0 – 19	5	5
20 – 39	11	16
40 – 59	26	
60 – 79	37	
80 – 99	9	
100 – 119	2	

(a) For the class interval 20 – 39, as written in the table above, complete the following sentences.

- (i) The upper class limit is
- (ii) The class width is
- (iii) Sixteen vehicles passed a checkpoint at no more thankmh⁻¹

SOLUTION:

Data: Cumulative frequency table showing the speeds, in kmh⁻¹, of 90 vehicles passing a certain checkpoint.

Required to complete: The sentences given for the class interval 20 – 39

Solution:

(See the modified table done)

- (i) The upper class limit is 39.
- (ii) The class width is $39.5 - 19.5 = 20$.
- (iii) Sixteen vehicles passed a checkpoint at no more than 39.5 kmh⁻¹.

(b) Complete the table shown above by inserting the missing values for the cumulative frequency column.

SOLUTION:

Required to complete: The cumulative frequency give.

Solution:

L.C.L-lower class limit, L.C.B-lower class boundary
U.C.L-upper class limit, U.C.B-upper class boundary


Speed, x , (in kmh ⁻¹)		Class Boundaries		Frequency	Cumulative Frequency	Points to Plot (U.C.B., CF)
L.C.L	U.C.L	L.C.B	U.C.B.			
						(0, 0)
0	19	$0 \leq x < 19.5$		5	5	(19.5, 5)
20	39	$19.5 \leq x < 39.5$		11	16	(39.5, 16)
40	59	$39.5 \leq x < 59.5$		26	$26 + 16 = 42$	(59.5, 42)
60	79	$59.5 \leq x < 79.5$		37	$37 + 42 = 79$	(79.5, 79)
80	99	$79.5 \leq x < 99.5$		9	$9 + 79 = 88$	(99.5, 88)
100	119	$99.5 \leq x < 119.5$		2	$2 + 88 = 90$	(119.5, 90)

- (c) On the grid provided, using a scale of 2 cm to represent 20 kmh^{-1} on the x – axis and 2 cm to represent 10 vehicles on the y – axis, draw the cumulative frequency curve to represent the information in the table.

SOLUTION:

Required to draw: The cumulative frequency curve to illustrate the information given

Solution:




- (d) (i) On your graph, draw reference lines to estimate the speed at which no more than 50% of the vehicles drove as they passed the check point.

SOLUTION:

Required to show: The speed at which no more than 50% of the vehicles drove past the checkpoint using reference lines

Solution:



- (ii) What is the estimated speed?


SOLUTION:

Required to state: The estimated speed at which no more than 50% of the vehicles passed the checkpoint

Solution:

The estimated speed is 62 kmh^{-1}

8. The first four figures in a sequence are shown below. Figure 1 is a single black dot, while each of the others consist of black dots arranged in an equilateral manner.




- (a) Draw Figure 5 of the sequence in the space below.

SOLUTION:

Data: Figures showing a sequence of black dots arranged in an equilateral manner.

Required to draw: The fifth figure in the sequence

Solution:



- (b) How many dots would be in Figure 6?

SOLUTION:

Required to state: The number of dots in Figure 6

Solution:

Figure	Number of Dots
1	1
2	3
3	6
4	10
5	15

By observation, the number of dots in the figure appear to be,
 $\frac{1}{2} \times (\text{the number of the figure}) \times (1 \text{ added to the number of the figure})$

$$\therefore \text{Figure 6 should have } \frac{1}{2} \times 6(6+1) = 21 \text{ dots}$$

The table below refers to the figures and the number of dots in each figure. Study the patterns below.

Figure, n	Number of Dots, d , in terms n	Number of Dots Used, d
1	$\frac{1}{2} \times 1 \times (1+1)$	1
2	$\frac{1}{2} \times 2 \times (2+1)$	3
3	$\frac{1}{2} \times 3 \times (3+1)$	6
11		
n		

- (c) Complete the row which corresponds to Figure 11 in the table above.

SOLUTION:

Data: Table showing the pattern of the number of dots used in the sequence of figures.

Required to complete: The row in the table that corresponds to Figure 11

Solution:

Figure, n	Number of Dots, d , in terms n	Number of Dots Used, d
1	$\frac{1}{2} \times 1 \times (1+1)$	1
2	$\frac{1}{2} \times 2 \times (2+1)$	3
3	$\frac{1}{2} \times 3 \times (3+1)$	6
11	$\frac{1}{2} \times 11 \times (11+1)$	66
n		

- (d) Determine which figure in the sequence has 210 dots.

SOLUTION:

Required to find: The figure that has 210 dots in the sequence

Solution:

$$\frac{1}{2}n(n+1) = 210$$

$$n(n+1) = 420$$

$$= 20 \times (20+1)$$

$$\therefore n = 20$$

\therefore Figure 20 has 210 dots.

- (e) Write a simplified algebraic expression for the number of dots, d , in the Figure n .

SOLUTION:

Required to state: An algebraic expression for the number of dots, d , in Figure n

Solution:

$$\frac{1}{2} \times n \times (n+1) = \frac{1}{2}n(n+1)$$

The completed table looks like:

Figure, n	Number of Dots, d , in terms n	Number of Dots Used, d
1	$\frac{1}{2} \times 1 \times (1+1)$	1
2	$\frac{1}{2} \times 2 \times (2+1)$	3
3	$\frac{1}{2} \times 3 \times (3+1)$	6
11	$\frac{1}{2} \times 11 \times (11+1)$	66
n	$\frac{1}{2} \times n \times (n+1)$	$\frac{1}{2} n(n+1)$

- (f) Show that there is no diagram that has exactly 1 000 dots.

SOLUTION:

Required To Show: No diagram in the sequence has 1 000 dots.

Solution:

$$\text{Let } \frac{1}{2}n(n+1) = 1000$$

$$n(n+1) = 2000$$


There are no two consecutive integers, n and $n+1$ being consecutive integers, whose product is exactly 2 000. Therefore, no diagram has 1 000 dots.

SECTION II

Answer TWO questions in this section.

ALGEBRA AND RELATIONS, FUNCTIONS AND GRAPHS

9. (a) The velocity – time graph below shows the motion of a cyclist over a period of 40 seconds.



- (i) Calculate the gradient of

- a) OA

SOLUTION:

Data: A velocity – time graph showing the motion of a cyclist for a 40 second period.

Required to find: The gradient of OA .

Solution:

$$O = (0, 0) \text{ and } A = (25, 10)$$

$$\text{Gradient of } OA = \frac{10 - 0}{25 - 0} = \frac{2}{5}$$

- b) AB

SOLUTION:

Required to find: The gradient of AB .

Solution:

$$A = (25, 10) \text{ and } B = (40, 10)$$

$$\text{Gradient of } AB = \frac{10 - 10}{40 - 25} = 0 \text{ (as expected for a horizontal line)}$$

- (ii) Complete the following statements.

The cyclist started from rest, where his velocity was ms^{-1} , and steadily increased his velocity by ms^{-1} each second during the first 25 seconds. During the next 15 seconds, his velocity remained constant, that is, his acceleration was ms^{-2} .

SOLUTION:

Required to complete: The sentences given.

Solution:

The cyclist started from rest, where his velocity was 0 ms^{-1} , and steadily increased his velocity by $\frac{2}{5} \text{ ms}^{-1}$ each second during the first 25 seconds.

During the next 15 seconds, his velocity remained constant, that is, his acceleration was 0 ms^{-2} .

- (iii) Determine the average speed of the cyclist over the 40-second period.

SOLUTION:

Required to find: Average speed of the cyclist over the 40 second period.

Solution:

$$\text{Average speed} = \frac{\text{Total distance covered}}{\text{Total time taken}}$$

$$\text{Area under the graph} = \text{Distance covered}$$

$$= \frac{1}{2} \{ (40 - 25) + (40 - 0) \} \times 10$$

$$= \frac{1}{2} (15 + 40) \times 10$$

$$= 275 \text{ m}$$

$$\therefore \text{Average speed} = \frac{275 \text{ m}}{40 \text{ s}}$$

$$= 6.875 \text{ ms}^{-1} \text{ or } 6\frac{7}{8} \text{ ms}^{-1}$$

- (b) Consider the following pair of simultaneous equations:

$$x^2 + 2xy = 5$$

$$x + y = 3$$

- (i) WITHOUT solving, show that $(1, 2)$ is a solution for the pair of simultaneous equations.

SOLUTION:

Data: The pair of simultaneous equations $x^2 + 2xy = 5$ and $x + y = 3$.

Required to show: $(1, 2)$ is a solution for the pair of simultaneous equations without solving

Solution:

Substitute $x = 1$ and $y = 2$ into both equations to test for ‘truth or falsity’

When $x = 1$ and $y = 2$

$$x^2 + 2xy = 5$$

$$(1)^2 + 2(1)(2) = 5$$

$$1 + 4 = 5$$

$$5 = 5 \text{ (True)}$$

When $x = 1$ and $y = 2$

$$x + y = 3$$

$$1 + 2 = 3$$

$$3 = 3 \text{ (True)}$$

$x = 1$ and $y = 2$ satisfy both equations

$\therefore (1, 2)$ is a solution for the pair of simultaneous equations.

- (ii) Solve the pair of simultaneous equations above to determine the **other** solution.

SOLUTION:

Required to find: The other solution for the pair of simultaneous equations.

Solution:

$$\text{Let } x^2 + 2xy = 5 \quad \dots \textcircled{1}$$

$$x + y = 3 \quad \dots \textcircled{2}$$

From equation $\textcircled{2}$

$$y = 3 - x \quad \dots \textcircled{3}$$

Substitute equation $\textcircled{3}$ into equation $\textcircled{1}$

$$x^2 + 2x(3 - x) = 5$$

$$x^2 + 6x - 2x^2 = 5$$

$$-x^2 + 6x - 5 = 0$$

$$\times -1$$

$$x^2 - 6x + 5 = 0$$

$$(x - 1)(x - 5) = 0$$

$$x - 1 = 0$$

$$x = 1$$

$$x - 5 = 0$$

$$x = 5$$

We already know that $x = 1$ is part of one set of solution. We consider the other.

$$\text{When } x = 5$$

$$y = 3 - x$$


$$= 3 - 5$$

$$= -2$$

\therefore The other solution is $(5, -2)$.

MEASUREMENT, GEOMETRY AND TRIGONOMETRY

10. (a) P, Q, R and S are four points on the circumference of the circle shown below. Angle $QRS = 58^\circ$.



Using the geometrical properties of a circle to give reasons for each step of your answer, determine the measure of


- (i) $\angle SPQ$

SOLUTION:

Data: Diagram of a circle where P, Q, R and S lie on its circumference and angle $QRS = 58^\circ$.

Required to calculate: $\angle SPQ$

Calculation:



$$\begin{aligned}\angle SPQ &= 180^\circ - 58^\circ \\ &= 122^\circ\end{aligned}$$


(The opposite angles of a cyclic quadrilateral are supplementary.)

(ii) $\angle OQS$

SOLUTION:


Required to calculate: $\angle OQS$

Calculation:



$$\begin{aligned}\angle SOQ &= 2(58^\circ) \\ &= 116^\circ\end{aligned}$$

(The angle subtended by a chord at the center of a circle is twice the angle that the chord subtends at the circumference, standing on the same arc.)




$\angle OSQ = \angle OQS$ (The base angles of an isosceles triangle are equal)

$$\begin{aligned}\angle OSQ + \angle OQS &= 180^\circ - 116^\circ \\ &= 64^\circ\end{aligned}$$

$$\begin{aligned}\therefore \angle OQS &= \frac{1}{2}(64^\circ) \\ &= 32^\circ\end{aligned}$$

- (b) A ship leaves Port A and sails 52 km on a bearing of 044° to Port B. The ship then changes course to sail to Port C, 72 km away, on a bearing of 105° .
- (i) On the diagram below, **not drawn to scale**, label the known distances travelled and the known angles.




SOLUTION:

Data: A ship leaves Port A and sails on a bearing of 044° to Port B, 52 km away. It then sails to Port C on a bearing of 105° , 72 km away.

Required to label: The sides and angles given on the diagram

Solution:




- (ii) Determine the measure of $\angle ABC$.

SOLUTION:

Required to find: $\angle ABC$

Solution: Let the line NB be extended to S, as shown.



$$\angle ABS = 44^\circ \quad (\text{Alternate angles are equal})$$

$$\begin{aligned} \angle CBS &= 180^\circ - 105^\circ \\ &= 75^\circ \end{aligned}$$

(two angles which make a straight line are supplementary)

$$\begin{aligned} \therefore \angle ABC &= 44^\circ + 75^\circ \\ &= 119^\circ \end{aligned}$$

- (iii) Calculate, to the nearest km, the distance AC.

SOLUTION:

Required to calculate: AC

Calculation:

By the cosine rule:


$$\begin{aligned} AC^2 &= (52)^2 + (72)^2 - 2(52)(72)\cos(119^\circ) \\ &= 2704 + 5184 - (-3630.25) \\ &= 11518.25 \\ AC &= \sqrt{11518.25} \\ &= 107.3 \text{ km} \\ &= 107 \text{ km (to the nearest km)} \end{aligned}$$

- (iv) Show that the bearing of A from C , to the nearest degree, is 260° .

SOLUTION:

Required to show: The bearing of A from C is 260°

Solution:



$$\angle BCN = 75^\circ \text{ (Co-interior angles are supplementary)}$$

By the sine rule:

$$\begin{aligned} \frac{52}{\sin \hat{A}CB} &= \frac{107.3}{\sin 119^\circ} \\ \sin \hat{A}CB &= \frac{52 \sin 119^\circ}{107.3} \\ &= 0.4238 \\ \hat{A}CB &= \sin^{-1}(0.4238) \\ &= 25.07^\circ \end{aligned}$$

$$\begin{aligned} \text{The bearing of } A \text{ from } C &= 360^\circ - (75^\circ + 25.07^\circ) = 259.9^\circ \\ &= 260^\circ \text{ (correct to the nearest degree)} \end{aligned}$$

Q.E.D.

VECTORS AND MATRICES

11. (a) Matrices A and B are such that

$$A = \begin{pmatrix} 3 & 2 \\ 5 & 4 \end{pmatrix} \text{ and } B = \begin{pmatrix} 4 & 0 \\ 3 & -1 \end{pmatrix}.$$

(i) Show by multiplying A and B , that $AB \neq BA$.

SOLUTION:

Data: $A = \begin{pmatrix} 3 & 2 \\ 5 & 4 \end{pmatrix}$ and $B = \begin{pmatrix} 4 & 0 \\ 3 & -1 \end{pmatrix}$

Required to show: $AB \neq BA$

Proof:

$$A \times B = \begin{pmatrix} 3 & 2 \\ 5 & 4 \end{pmatrix} \begin{pmatrix} 4 & 0 \\ 3 & -1 \end{pmatrix}$$

$2 \times 2 \times 2 \times 2 = 2 \times 2$

$$= \begin{pmatrix} e_{11} & e_{12} \\ e_{21} & e_{22} \end{pmatrix}$$

$$\begin{aligned} e_{11} &= (3 \times 4) + (2 \times 3) \\ &= 12 + 6 \\ &= 18 \end{aligned}$$

$$B \times A = \begin{pmatrix} 4 & 0 \\ 3 & -1 \end{pmatrix} \begin{pmatrix} 3 & 2 \\ 5 & 4 \end{pmatrix}$$

$2 \times 2 \times 2 \times 2 = 2 \times 2$

$$= \begin{pmatrix} e_{11} & e_{12} \\ e_{21} & e_{22} \end{pmatrix}$$

$$\begin{aligned} e_{11} &= (4 \times 3) + (0 \times 5) \\ &= 12 + 0 \\ &= 12 \end{aligned}$$

e_{11} of $AB \neq e_{11}$ of BA . There is no need for further working since, regardless of whether the remaining three corresponding entries of AB are equal or not to those of BA , all the entries of AB will never be equal to the entries of BA . $\therefore AB \neq BA$.

Q.E.D.

(ii) Find A^{-1} , the inverse of A .

SOLUTION:

Required to find: A^{-1}

Solution:

$$\begin{aligned}\det A &= (3 \times 4) - (2 \times 5) \\ &= 12 - 10 \\ &= 2\end{aligned}$$

$$\begin{aligned}A^{-1} &= \frac{1}{2} \begin{pmatrix} 4 & -(2) \\ -(5) & 3 \end{pmatrix} \\ &= \begin{pmatrix} 2 & -1 \\ -\frac{5}{2} & \frac{3}{2} \end{pmatrix}\end{aligned}$$

- (iii) Write down the 2×2 matrix representing the matrix product AA^{-1} .

SOLUTION:

Required to write: The 2×2 matrix AA^{-1} .

Solution:

$$\begin{aligned}A \times A^{-1} &= I \\ 2 \times 2 \times 2 \times 2 &= 2 \times 2\end{aligned}$$

$$I = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

- (b) (i) Write the following pair of simultaneous equations as a matrix equation.

$$\begin{aligned}3x + 2y &= 1 \\ 5x + 4y &= 5\end{aligned}$$

SOLUTION:

Data: $3x + 2y = 1$ and $5x + 4y = 5$

Required to write: The given pair of simultaneous equations as a matrix equation

Solution:

$$\begin{pmatrix} 3 & 2 \\ 5 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

- (ii) Write the solution of your matrix equation in (b) (i) as a product of two matrices.

SOLUTION:

Required to write: The solution of the matrix equation as a product of two matrices

Solution:

$$\begin{pmatrix} 3 & 2 \\ 5 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$A \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$\times A^{-1}$

$$A \times A^{-1} \begin{pmatrix} x \\ y \end{pmatrix} = A^{-1} \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$


$$I \begin{pmatrix} x \\ y \end{pmatrix} = A^{-1} \begin{pmatrix} 1 \\ 5 \end{pmatrix}$$

$$\begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 2 & -1 \\ -\frac{5}{2} & \frac{3}{2} \end{pmatrix} \begin{pmatrix} 1 \\ 5 \end{pmatrix} \text{ as a product of two matrices.}$$

If calculated, (which was not required according to the question) the right hand side would give $\begin{pmatrix} -3 \\ 5 \end{pmatrix}$ and equating corresponding entries gives $x = -3$ and $y = 5$.

- (c) The position vectors of the points P and Q relative to an origin O , are $OP = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$ and $OQ = \begin{pmatrix} 5 \\ 0 \end{pmatrix}$ respectively.

The diagram below shows that $PR = 3OP$ and $QS = 3OQ$.



(i) Express in the form $\begin{pmatrix} x \\ y \end{pmatrix}$, vector

- OS

SOLUTION:

Data: Diagram showing position of vectors of the points P and Q , where $OP = \begin{pmatrix} 4 \\ 3 \end{pmatrix}$ and $OQ = \begin{pmatrix} 5 \\ 0 \end{pmatrix}$ and $PR = 3OP$ and $QS = 3OQ$.

Required to express: OS in the form $\begin{pmatrix} x \\ y \end{pmatrix}$.

Solution:

$$OQ = \begin{pmatrix} 5 \\ 0 \end{pmatrix}$$

$$QS = 3 \begin{pmatrix} 5 \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} 15 \\ 0 \end{pmatrix}$$

$$OS = OQ + QS$$

$$= \begin{pmatrix} 5 \\ 0 \end{pmatrix} + \begin{pmatrix} 15 \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} 20 \\ 0 \end{pmatrix} \text{ and is of the form } \begin{pmatrix} x \\ y \end{pmatrix}, \text{ where } x = 20 \text{ and } y = 0.$$

- PQ

SOLUTION:

Required to express: PQ in the form $\begin{pmatrix} x \\ y \end{pmatrix}$.

Solution:

$$PQ = PO + OQ$$

$$= - \begin{pmatrix} 4 \\ 3 \end{pmatrix} + \begin{pmatrix} 5 \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} 1 \\ -3 \end{pmatrix} \text{ and is of the form } \begin{pmatrix} x \\ y \end{pmatrix}, \text{ where } x = 1 \text{ and } y = -3.$$

- RS

SOLUTION:

Required to express: RS in the form $\begin{pmatrix} x \\ y \end{pmatrix}$.

Solution:

$$PR = 3OP$$

$$= 3 \begin{pmatrix} 4 \\ 3 \end{pmatrix}$$

$$= \begin{pmatrix} 12 \\ 9 \end{pmatrix}$$

$$OR = OP + PR$$

$$= \begin{pmatrix} 4 \\ 3 \end{pmatrix} + \begin{pmatrix} 12 \\ 9 \end{pmatrix}$$

$$= \begin{pmatrix} 16 \\ 12 \end{pmatrix}$$

$$RS = RO + OS$$

$$= - \begin{pmatrix} 16 \\ 12 \end{pmatrix} + \begin{pmatrix} 20 \\ 0 \end{pmatrix}$$

$$= \begin{pmatrix} 4 \\ -12 \end{pmatrix} \text{ and is of the form } \begin{pmatrix} x \\ y \end{pmatrix}, \text{ where } x = 4 \text{ and } y = -12.$$

- (ii) State TWO geometrical relationships between PQ and RS .

SOLUTION:

Required to state: Two geometrical relationships between PQ and RS .

Solution:

$$PQ = \begin{pmatrix} 1 \\ -3 \end{pmatrix}$$

$$RS = \begin{pmatrix} 4 \\ -12 \end{pmatrix}$$

$$= 4 \begin{pmatrix} 1 \\ -3 \end{pmatrix}$$

$$= 4PQ$$

Hence, RS is a scalar multiple (which is 4) of PQ . Therefore, RS and PQ are parallel.

$$RS = 4 \begin{pmatrix} -1 \\ 3 \end{pmatrix}$$

$$= 4 \times PQ$$

$$|RS| = 4|PQ|$$

That is to say, the length of RS is 4 times the length of PQ .

www.faspassmaths.com